

# Time-resolved biophysical methods in the study of protein folding

Kevin W Plaxco\* and Christopher M Dobson†

Many of the biophysical techniques developed to characterize native proteins at equilibrium have now been adapted to the structural and thermodynamic characterization of transient intermediate populations during protein folding. Recent advances in these techniques, the use of novel methods of initiating refolding, and a convergence of theoretical and experimental approaches are leading to a detailed understanding of many aspects of the folding process.

## Addresses

New Chemistry Laboratory, University of Oxford, South Parks Road, Oxford OX1 3QT, UK

\*e-mail: kwp@bioch.ox.ac.uk

†e-mail: chris.dobson@icl.ox.ac.uk

Current Opinion in Structural Biology 1996, 6:630–636

© Current Biology Ltd ISSN 0959-440X

## Abbreviations

<b>ANS</b>	8-anilino-1-naphthalenesulphonate
<b>C12</b>	chymotrypsin inhibitor 2
<b>MS</b>	mass spectrometry
<b>NOE</b>	nuclear Overhauser effect
<b>SAXS</b>	small-angle X-ray scattering
<b>T-jump</b>	temperature jump

## Introduction

The primary question addressed in studies of protein folding can be stated very simply: how do denatured polypeptide chains limit their conformational search in order to achieve the native state in a biologically relevant time? The complexity of the denatured state rules out the possibility that folding is a simple stochastic search process [1], and folding is, almost certainly, facilitated by the existence of (potentially multiple) specific mechanisms. Much effort has gone into the characterization of transient partially folded states arising during folding [2,3•] in an attempt to understand these mechanisms and the process by which proteins rapidly fold to their native structures.

Native proteins are characterized by a high degree of compactness, an ordered hydrophobic core, a well defined overall architecture, and the presence of specific and cooperative interactions among buried side chains. Recent progress in both instrumentation and experimental design has provided unprecedented insights into the evolution of each of these characteristics as an initially disordered and extended polypeptide chain folds via a heterogeneous population of partially folded states into its native conformation (Fig. 1). In this article we focus on recent advances in the time-resolved experimental characterization of the properties and distribution of

partially folded states arising during nonoxidative refolding *in vitro*, and the promise that these developments hold for providing a detailed description of the folding process.


## The initiation of folding

Protein folding in the cell follows synthesis of the polypeptide chain on a ribosome. Refolding *in vitro* is more readily initiated by rapidly transferring a protein from denaturing conditions to an environment in which the native conformation is favoured. This is often achieved by diluting protein solutions containing denaturant with nondenaturing buffers using a stopped-flow mixing device. Turbulent mixers, such as the Berger ball mixer used in many commercially available instruments, achieve high mixing efficiency by interweaving fine, turbulence-generated streams [4]. The minimum dimensions of these streams is limited by technical issues, such as cavitation, such that denaturants require >100 μs to diffuse from them. Limits on the physical proximity of a detecting cell to the mixer, and the speed with which flow can be stopped without producing shock effects, further increase the deadtime of most stopped-flow instruments to >1 ms.

The extremely rapid burst-phase events now evident for many proteins are complete within the deadtime of conventional stopped-flow mixing devices [5]. Fortunately, recent technical advances promise significant reductions in these mixing deadtimes. The use of continuous-flow devices that avoid the shock disturbances of high-speed stopped-flow and 'freejet' mixers, which generate small, rapidly diffusing streams by laminar flow through very small orifices, has lowered deadtimes to tens of microseconds [6•]. Non-mixing methods, such as flash photolysis [7], optical electron transfer [8•] and temperature jump (T-jump) [9•,10•], promise further improvements. Optical electron transfer, based on the existence of conditions under which an oxidized redox protein is unfolded but the reduced form is native, has been used to initiate the refolding of cytochrome *c* in <1 μs by photochemically induced reduction [8•]. T-jump experiments, based on reversing cold-induced denaturation through rapid sample heating, have yielded deadtimes of ~10 μs by electrical-discharge heating [9•] and an amazing ~20 ns by laser-induced heating [10•]. Applied to the folding of apomyoglobin, laser T-jump has been used to characterize a collapsed state formed in a diffusion-limited reaction that is completed within ~20 μs [10•]. When coupled with high-speed absorbance, fluorescence and CD, these new folding-initiation techniques will undoubtedly provide important insights into the chemistry of the earliest events in folding.

**Figure 1**

A schematic representation of the characteristics of globular proteins that can be followed during refolding, with time resolution in the second to millisecond range. Other properties that can be monitored, but are not indicated, include the creation and disruption of organized hydrophobic voids and overall thermodynamic stability. Although no individual probe can monitor all of the structural details of a folding intermediate, the use of multiple complementary probes can provide a detailed picture of the distribution of conformations that make up transient folding populations.



### Measuring collapse and core packing

A general property of protein folding is that an extended and highly disordered polymer chain must collapse to form a compact, globular protein [11•]. Measures of molecular dimensions and core packing (Table 1) are thus critical elements of a complete description of the folding process. Indirect probes of these properties, such as changes in the UV absorbance of aromatic residues [12•], the fluorescence of tryptophan or tyrosine side chains [13••], or the fluorescence of extrinsic fluorophores such as 8-anilino-1-naphthalenesulphonate (ANS) [14•], have seen widespread application. More direct probes of the exclusion of solvent from the hydrophobic core, involving monitoring the accessibility of hydrophilic fluorescence quenchers such as iodide or acrylamide [13••] or the reactivity of cysteine side chains [15], have also been developed. The use of time-resolved fluorescence spectroscopy, not only to monitor molecular dimensions but also to provide a detailed description of the loss of core residue mobility during the refolding of dihydrofolate reductase [16•], is a recent example of the variety of indirect indicators of the collapse and core packing that are available. What most of these probes lack, however, is an ability to monitor the distribution of individual species in heterogeneous mixtures or to provide a quantitative measure of the dimensions of partially folded conformations. Although little progress has

been made on the former, several quantitative probes of molecular dimension are now available.

Time-resolved fluorescence energy transfer, small-angle X-ray scattering (SAXS), and quasi-elastic light scattering have all been used to provide a direct measurement of the dimensions of species arising during folding. The detection of fluorescence energy transfer between a covalently attached fluorophore and a tryptophan side chain, which has been used to attempt direct measurements of the evolution of collapsed species during the refolding of apomyoglobin [17•] and other proteins [18], is consistent with the hypothesis that these proteins fold via a rapidly formed intermediate of near-native compactness. Such studies are, however, limited to proteins that can be modified with suitable fluorophores and only provide measurements of a single scalar distance. Unlike fluorescence energy transfer, SAXS [19] and quasi-elastic light scattering provide direct means of monitoring the overall dimensions of macromolecules. SAXS, when implemented with very high flux synchrotron X-ray sources, provides a measure of the average radius of gyration with <100 ms time resolution. This technique has recently been applied to the refolding of apomyoglobin, again indicating the near-native compactness of the major folding intermediate of this protein [20•,21]. Quasi-elastic light scattering, though presently limited by a ~1 s deadtime, monitors

Table 1

**Biophysical techniques used to investigate protein folding\*.**

Property	Technique	Resolution	Measurement	Reference
Core packing	Intrinsic fluorescence	< 1 ms	The orientation and environment of (predominantly) tryptophan side chains	[13**]
	Ultraviolet absorbance	ms	The orientation and environment of (predominantly) tyrosine side chains	[12*]
	Extrinsic (ANS) fluorescence	ms	Formation and disruption of organized hydrophobic patches and clefts	[14*]
	Fluorescence quenching	ms	Isolation of tryptophan side chains from hydrophilic fluorescence quenchers	[13**]
	CysteinyI quenching	10 s	Protection of cysteine side chains from hydrophilic reactants	[15]
Molecular dimensions	Fluorescence anisotropy	ms	Tryptophan side chain mobility and overall molecular dimensions	[16*]
	Fluorescence energy transfer	ms	Scalar distance between tryptophan and a covalently attached fluorophore	[17*]
	Small angle X-ray scattering	< 100 ms	The average radius of gyration	[20*]
	Quasi-elastic light scattering	1 s	The average radius of gyration	[22]
Secondary structure and persistent hydrogen bonds	Far-UV circular dichroism	ms	Backbone conformation averaged over sequence and population	[5]
	Pulse labelling NMR	5–10 ms	Sequence specific formation of stable amide and tryptophan hydrogen bonds	[24]
	Pulse labelling mass spectrometry	5–10 ms	The formation of persistent hydrogen bonds in discrete intermediates	[25*]
Tertiary contacts and native structure	Biological activity	ms–s	The formation of native tertiary structure at the active site	[5]
	Interrupted folding	10 ms	The unfolding rate of discrete intermediates as a probe of their stability	[30]
	Near-UV circular dichroism	ms	Formation of stable aromatic and disulphide bond tertiary contacts	[5]
	Real-time NMR	1 s	Formation of specific side chain tertiary contacts	[35**]
	Protein engineering	†	The energetic contributions of side chains to discrete intermediates	[32**]

\*Many of the biophysical techniques developed to characterize native proteins at equilibrium have now been adapted to the structural and thermodynamic characterization of transient populations during folding. Here we summarize many of the biophysical techniques that have been used in recent years to characterize the folding of a variety of proteins. A single reference to each method is provided that either reflects a recent review of the subject or an illustrative application of the technique. †The time resolution of protein-engineering refolding experiments is limited only by the time resolution of the probe used to monitor folding mutants.

the translational mobility and thus overall dimensions of a macromolecule, and has been used to probe the formation of compact states during the refolding of lysozyme [22]. There appears to be no fundamental reason why these techniques will not prove to be general methods for observing directly the dimensions of a polypeptide chain during protein folding.

### Monitoring the formation of secondary structure

Probes of the backbone conformation, such as far-UV CD and pulse-labelling hydrogen exchange, have provided a wealth of data on the kinetics of secondary-structure formation during folding (Table 1). The recovery of far-UV CD ellipticity is widely considered a critical measure of the average secondary structure content in heterogeneous folding mixtures. For proteins, however, much of the formation of secondary structure occurs in a burst phase during the mixing deadtime and thus has not been amenable to direct study. It is fortunate,

then, that the invention of extremely rapid methods for the initiation of refolding comes close on the heels of advances in high-speed CD [23]. Now that the application of high-intensity laser light sources to CD spectropolarimetry has produced sub- $\mu$ s time resolution, fundamental questions about the timing of the formation of secondary structure may soon be answered.

Although CD provides an estimate of average secondary structure content, it does not provide information on the specific residues involved or the distribution of conformations present. Pulse-labelling amide-exchange experiments can provide this complementary information by monitoring the formation of stable backbone hydrogen bonds [24]. Pulse labelling linked to NMR spectroscopy has been used for a number of years as a probe of the sequence-specific formation of persistent elements of secondary structure but, like optical methods, the technique cannot resolve individual components from heterogeneous mixtures. Advances in coupling pulse

labelling and mass spectrometry (MS) have furthered our understanding of the formation of secondary structure by allowing the observation of resolved molecular species. This has provided a means of characterizing the hydrogen-exchange properties of discrete species in heterogeneous populations, as observed, for example, during the refolding of lysozyme [25•]. MS, like optical methods, provides data averaged over the entire sequence of a molecule. Technical advances in MS, however, have proven the feasibility of identifying the sequences of protein cleavage products produced in the gas phase by collision-induced dissociation [26,27•]. It may thus soon prove possible to produce sequence-specific hydrogen-exchange data for discrete species in complex folding populations.

### Detecting tertiary contacts

Because the formation of partially ordered states with regions of native-like structure is thought to be an essential step in protein folding, detecting native tertiary contacts in transient folding populations has been a major goal of folding research. Near-UV CD, which primarily monitors the aromatic side chains immobilized by asymmetric tertiary contacts, has proved an important probe of the recovery of native structure [5]. For many proteins, time-resolved assays of the recovery of biological activity (e.g. the binding of fluorescent substrates or inhibitors) can be used to monitor the recovery of a native active site [12•,28]. Interrupted folding experiments, in which transiently refolded mixtures undergo a second unfolding by the rapid addition of denaturant, have been used to detect the formation of material with native stability [29]. This method, which relies on the reasonable assumption that the unfolding rate of a given conformation reflects its thermodynamic stability, has recently been used to monitor the stability of an intermediate in the folding of barnase [30] and to support the existence of parallel pathways in the folding of lysozyme [31•]. It may provide a general probe of the formation of both native and near-native structures.

Although probes of the formation of native protein are well established, only recently have techniques been developed that can monitor the formation of specific tertiary contacts during folding. Protein engineering provides one method of assaying the contributions of specific side chains to the energetics of transient folding intermediates. The contributions of these side chains (relative to their contribution to the stability of the native protein) have been interpreted as a measure of the 'nativeness' of their contacts in the intermediate. Major folding intermediates of barnase [32••] and phosphoglycerate kinase [33•] have been characterized using this technique. Stopped-flow NMR has also been used to monitor the formation of specific tertiary interactions during folding [34,35••]. Though presently limited to a time resolution of ~1 s, the technique provides a nonperturbing method of detecting the formation of the highly shifted resonances

characteristic of native proteins. This method shows much potential for providing information concerning the formation of specific native and native-like contacts during folding.

### Transition-state probes

A complete description of the folding process requires knowledge of both the structure and energetics of the rate-determining conformation. While the ephemeral nature of transition states generally precludes direct structural studies, the transition state is the conformation of the rate-limiting step (or steps) and therefore the kinetics of folding can provide an indirect probe of its structure. The effect of environmental factors and mutations on the kinetics of the recovery of native properties (such as fluorescence) have thus been used to provide a detailed picture of the conformation of this most fleetingly transient species in protein folding.

Environmental factors that affect folding rates have provided valuable clues to the general nature of folding transition states. For example, studies of the temperature dependence of protein folding rates have been used to probe their thermodynamic properties [36•]. Other studies into the effects of pressure [37•], denaturants [38], ionic strength [39] and pH [40], have been used to define relative molar volumes and solvent-exposed surface areas of transition states, and to probe the contributions of ionizable groups to their energetics. From such studies, a general picture is emerging of a typical transition state as a collapsed but still relatively poorly packed set of conformations.

Efforts to ascertain the high-resolution structure of a folding transition state have focused on protein engineering experiments designed to produce a map of the energetic contributions of specific side chains to the rate-limiting step. This has been carried out in some detail for barnase and chymotrypsin inhibitor 2 (CI2) [32••] in studies that have provided insights into the structure and heterogeneity of the transition state [41], and suggest that small nuclei of native-like structure are involved in the rate-determining steps of the folding of at least some proteins.

### Conclusions

As the number and quality of biophysical techniques with sufficient time resolution increases, so our detailed knowledge of the folding process improves. Issues such as cooperativity, collapse and the formation of secondary structure during refolding are becoming well described for a number of proteins. What is still lacking, however, is a means of generating a picture of the structure and distributions of transient folding populations with good spatial resolution. The next challenge in protein folding lies in discovering how to produce such high-resolution data. Several potential approaches now appear feasible.

Because no single method can provide a complete picture of the distribution of structures in a transient population of intermediates (Fig. 1), it is clear that multiple complementary approaches must be combined to generate detailed structural models. For example, dynamic light-scattering and intrinsic fluorescence can be used to define the average dimensions and degree of core packing in a population, pulse-labelling amide exchange can provide information on the location and stability of secondary structure, and NMR and inhibitor binding can be used to define specific tertiary contacts. Such information can thus be brought together to develop a detailed picture of the key features the folding process [12•,42•,43•].

The direct acquisition of high-resolution structural information may also be possible through modifications of current biophysical methods. NMR, for example, was converted from a technique of low spatial resolution to one applicable to high spatial resolution by the introduction of two-dimensional spectroscopy. The application of multi-dimensional NMR techniques to the study of highly transient structures may appear daunting, but the increased availability of specifically isotopically labeled proteins and high-field spectrometers has already made possible two-dimensional refolding experiments with a time resolution of a few minutes [44•] by repetitive accumulation of rapidly acquired spectra. Novel approaches may provide significant further reductions. For example, the refolding of a protein during the acquisition of two-dimensional data can result in changes in peak shape that can be deconvoluted to provide a wealth of information on the refolding kinetics of individual elements of the protein with time resolution on the order of seconds. Further potential exists for experiments in which nuclear Overhauser effect (NOE) crosspeaks generated in folding populations are detected in the spectrally well characterized native protein, possibly to provide a detailed picture of the tertiary contacts formed in transient folding intermediates (J Balbach *et al.*, unpublished data).

Theoretical methods can provide atomic-level models of the structure and distribution of protein folding intermediates but they necessarily involve significant simplifying assumptions. The iterative coupling of simulation with experiment may provide the necessary constraints on these assumptions to produce accurate high-resolution models. Recent studies of the details of denatured protein conformations have provided an example of this type of complementary theoretical and experimental approach. Current understanding of the 'random coil' denatured state has recently been advanced by the use of experiments to verify the specific predictions produced from Monte Carlo simulations of the denatured state. Further refinements of these simulations based on discrepancies between predicted and observed NOEs and J-coupling

constants [45••] has led to a deeper understanding of the conformational distributions within denatured states.

A similar combined simulation and experimental approach has been used for the interpretation of protein engineering investigations into the structure of the folding transition states of CI2 and barnase. Molecular dynamics simulations, here inspired and constrained by experimental investigations, proved vital for the formation of high-resolution models of these folding transition states [32••,46••,47••]. The complementary aspects of theory and experiment in protein folding suggest that this promising trend will continue. In particular, it is to be hoped that, with the addition of experimentally derived constraints, lattice simulations [47••,48•,49•] will lead to higher-resolution models of intermediate populations with significant predictive value. The use of complementary biophysical approaches to obtain adequate information to constrain theoretical models holds great promise for providing a detailed description and understanding of the folding process.

## Acknowledgements

We thank Elaine Marzluff, Jay Winkler, Yuji Goto and members of the Dobson research group for generously sharing their expertise. This research is supported in part by an International Research Scholars award from the Howard Hughes Medical Institute to CM Dobson. The Oxford Centre for Molecular Sciences is supported by the UK Biotechnology and Biological Sciences Research Council, the Medical Research Council and the Engineering and Physical Sciences Research Council.

## References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Levinthal C: **Are there pathways for protein folding?** *J Chem Phys* 1968, **65**:44–45.
2. Baldwin RL: **Finding intermediates in protein folding.** *Bioessays* 1994, **16**:207–210.
3. Ptitsyn OB: **Structures of folding intermediates.** *Curr Opin Struct Biol* 1995, **5**:74–78.
- A discussion of the characteristics of both kinetic and equilibrium partially folded states, and the nature of the barriers that separate these conformations from denatured and fully native species.
4. Berger RL, Backo B, Chapman HF: **High resolution mixer for the study of rapid reactions in solution.** *Rev Sci Instr* 1968, **39**:493–498.
5. Evans PA, Radford SE: **Probing the structure of folding intermediates.** *Curr Opin Struct Biol* 1994, **4**:100–106.
6. Takahashi S, Ching Y-L, Wang J, Rousseau DL: **Microsecond generation of oxygen-bound cytochrome c oxidase by rapid solution mixing.** *J Biol Chem* 1995, **270**:8405–8407.
- A recent (but nonfolding) example of a continuous-flow freejet mixing device with a deadline in the tens of microseconds range. The use of a laser light source and continuous flow, which provides long signal integration times, highlights the requirement for very sensitive detection methods to achieve good time resolution.
7. Jones CM, Henry ER, Hu U, Chan C-K, Luck SD, Bhuyan A, Roder H, Hofrichter J, Eaton WA: **Fast events in protein folding initiated by nanosecond laser photolysis.** *Proc Natl Acad Sci USA* 1993, **90**:11860–11864.
8. Pascher T, Chesick JP, Winkler JR, Gray HB: **Protein folding triggered by electron transfer.** *Science* 1996, **271**:1558–1560.
-

The optically triggered reduction of oxidized, denatured cytochrome *c* initiates refolding in less than one microsecond. Time-resolved heme absorbance measurements suggest a rapid collapse (time constant 40  $\mu$ s in 4.6 M guanidine hydrochloride at 40°C) followed by slow progression to the native conformation.

9. Nöling B, Golbik R, Fersht AR: **Submillisecond events in protein folding.** *Proc Natl Acad Sci USA* 1995, **92**:10668–10672.

The refolding of barstar is initiated in ~10  $\mu$ s by rapidly heating the cold-denatured protein. Tryptophan fluorescence is used to identify a rapid phase (time constant 320  $\mu$ s at 10°C) that is attributable to the formation of a relatively solvent exposed and unconsolidated folding intermediate.

10. Ballew RM, Sabelko J, Gruebele M: **Direct observation of fast protein folding: the initial collapse of apomyoglobin.** *Proc Natl Acad Sci USA* 1996, **93**:5759–5764.

The laser-induced T-jump initiation of refolding is used to probe the extremely rapid initial stages of the refolding of apomyoglobin. Analysis of fluorescence changes upon initiation with ~15 ns time resolution indicates that a partially solvent-excluded collapsed state is achieved in a rapid reaction with a time constant of ~7  $\mu$ s at 22°C. The solvent viscosity dependence of this rate suggests that the formation of this state is limited by large-scale protein motions during the initial collapse.

11. Miranker AD, Dobson CM: **Collapse and cooperativity in protein folding.** *Curr Opin Struct Biol* 1996, **6**:31–42.

An overview of the evidence for and implications of the occurrence of a highly cooperative collapse during the refolding of globular proteins.

12. Udgaonkar JB, Baldwin RL: **Nature of the early folding intermediate of ribonuclease A.** *Biochemistry* 1995, **34**:4088–4096.

A variety of discrete intermediates have been described on the folding pathway of RNase A, in part on the basis of the ultraviolet absorbance characteristics of tyrosine side chains. In this paper, the properties of an early folding intermediate of RNase A are characterized by tyrosine absorbance, inhibitor binding (also monitored by absorbance) and pulse-labelling hydrogen exchange.

13. Engelhard M, Evans PA: **Experimental investigation of side chain interactions in early folding intermediates.** *Fold Des* 1996, **1**:31–37.

A comprehensive and up-to-date review of techniques available to monitor the evolution of side chain interactions during refolding, as a measure of both hydrophobic core organization and overall collapse of the polypeptide chain.

14. Engelhard M, Evans PA: **Kinetics of interaction of partially folded proteins with a hydrophobic dye: evidence that molten globule character is maximal in early folding intermediates.** *Protein Sci* 1996, **4**:1553–1562.

Binding of the hydrophobic dye ANS is considered the hallmark of the formation of a class of collapsed and loosely packed intermediates termed molten globules. In this paper, ANS is used in a carefully controlled manner to probe the evolution of this species during the folding of several proteins. A significant perturbative effect is noted, and a manner of avoiding this pitfall demonstrated.

15. Ballery N, Desmadril M, Minard P, Yon JM: **Characterization of an intermediate in the folding pathway of phosphoglycerate kinase: chemical reactivity of genetically introduced cysteinyl residues during the folding process.** *Biochemistry* 1993, **33**:708–714.

16. Jones BE, Beechem JM, Matthews CR: **Local and global dynamics during the folding of *E. coli* dihydrofolate reductase by time-resolved fluorescence spectroscopy.** *Biochemistry* 1995, **34**:1867–1877.

Time-resolved fluorescence spectroscopy provides detailed information on the dynamics of tryptophan side chains and the bound reporter group ANS. This study of the refolding of dihydrofolate reductase suggests that an early intermediate, despite being collapsed to a near-native radius of gyration, maintains side chain mobilities similar to those of the denatured state.

17. Rischel C, Poulsen FM: **Modification of a specific tyrosine enables tracing of the end-to-end distance during apomyoglobin folding.** *FEBS Lett* 1995, **374**:105–109.

An excellent example of data derived from fluorescence energy transfer experiments. Here, the normally vexing requirement of the covalent attachment of a suitable fluorophore is overcome by the chemical modification of a tyrosine side chain *in situ*. This study, like the SAXS studies discussed in [20\*], indicates that a collapsed intermediate with near-native separation of a specific tyrosine and tryptophan pair occurs early during the refolding of apomyoglobin.

18. Kawata Y, Hamaguchi K: **Use of fluorescence energy transfer to characterize the compactness of the constant fragment of an immunoglobulin light chain in the early stage of folding.** *Biochemistry* 1991, **30**:4367–4373.

19. Lattman EE: **Small angle scattering studies of protein folding.** *Curr Opin Struct Biol* 1994, **4**:87–92.

20. Eliezer D, Jennings PA, Wright PE, Doniach S, Hodgson SK, Tsuruta H: **The radius of gyration of an apomyoglobin folding intermediate.** *Science* 1995, **270**:487–488.

Direct determination of the radius of gyration of a putative molten-globule intermediate during the refolding of apomyoglobin with <100 ms time resolution. The value obtained suggests that apomyoglobin folds via a rapidly formed intermediate of near-native compactness.

21. Eliezer D, Chiba K, Tsuruta H, Doniach S, Hodgson KO, Kihara H: **Evidence of an associative intermediate on the myoglobin refolding pathway.** *Biophys J* 1995, **65**:912–917.

22. Feng H-P, Widom J: **Kinetics of compaction during lysozyme refolding studied by continuous-flow quasielastic light scattering.** *Biochemistry* 1994, **33**:13382–13390.

23. Zhang C-F, Lewis JW, Cerpa R, Kuntz ID, Kliger D: **Nanosecond circular dichroism spectral measurements: extension to the far-ultra violet range.** *J Phys Chem* 1993, **97**:5499–5505.

24. Baldwin RL: **Pulsed H/D-exchange studies of folding intermediates.** *Curr Opin Struct Biol* 1993, **3**:84–91.

25. Miranker A, Robinson CV, Radford SE, Dobson CM: **Investigation of protein folding by mass spectrometry.** *FASEB J* 1996, **10**:93–101.

A review of recent developments in the analysis of equilibrium and pulse-labelling hydrogen-exchange data by MS. Particular attention is paid to the ability of these techniques to resolve individual components from heterogeneous folding populations.

26. Anderegg RJ, Wagner DS, Stevenson CL, Borchardt RT: **The mass spectrometry of helical unfolding in peptides.** *J Am Soc Mass Spectrom* 1994, **5**:425–433.

27. Miranker A, Kruppa GH, Robinson CV, Aplin RT, Dobson CM: **An isotope labelling strategy for the assignment of protein fragments generated for mass spectrometry.** *J Am Chem Soc* 1996, in press.

This paper reports the use of isotope labelled proteins to aid in the identification of lysozyme fragments produced by chemically induced dissociation. This technique makes possible the identification of many of the small peptides produced by gas-phase dissociation and suggests that MS-derived sequence-specific hydrogen-protection studies may be possible.

28. Jennings PA, Finn BE, Jones BE, Mathews CR: **A reexamination of the folding mechanism of dihydrofolate reductase from *Escherichia coli*: verification and refinement of a four-channel model.** *Biochemistry* 1993, **32**:3783–3789.

29. Schmid FX: **Mechanism of folding of ribonuclease A. Slow refolding is a sequential reaction via structural intermediates.** *Biochemistry* 1983, **22**:4690–4696.

30. Schreiber G, Fersht AR: **The refolding of *cis*- and *trans*-peptidylprolyl isomers of barstar.** *Biochemistry* 1993, **32**:11195–11203.

31. Kiefhaber T: **Kinetic traps in lysozyme folding.** *Biochemistry* 1995, **92**:9029–9033.

The refolding of lysozyme is monitored with 10 ms time resolution (and a 20 ms deadtime) using an interrupted folding technique, in which transiently refolded protein is re-unfolded by the addition of denaturant. The rate of formation of native material is quantified by assuming that the fraction of protein exhibiting a slow unfolding rate represents native protein. The observation of a biphasic rate of recovery for this slowly unfolding material supports the view that this protein refolds via at least two parallel pathways.

32. Fersht AR: **Mapping the structures of transition states and intermediates in folding: delineation of pathways at high resolution.** *Philos Trans R Soc Lond* 1995, **348**:11–15.

A review of the application of site-directed mutagenesis techniques to the exploration of protein folding. Particular attention is paid to the characterization of the folding transition states of barnase and Cl2.

33. Parker MJ, Sessions RB, Badcoe IG, Clarke AR: **The development of tertiary interactions during the folding of a large protein.** *Fold Des* 1996, **1**:145–156.

Protein engineering is used to probe the formation of native contacts by the effect of mutations on the stability of a transient intermediate during the refolding of phosphoglycerate kinase.

34. Koide S, Dyson HJ, Wright PE: **Characterization of a folding intermediate of apoplastocyanin trapped by proline isomerization.** *Biochemistry* 1993, **32**:12299–12310.

35. Balbach J, Forge V, VanNuland NAJ, Winder SL, Hore PJ, Dobson CM: **Following protein-folding in real-time using NMR-spectroscopy.** *Nat Struct Biol* 1995, **2**:865–870.

The application of stopped flow NMR to the refolding of apo-lactalbumin demonstrates the existence of a rapidly formed kinetic molten-globule intermediate previously postulated from extrinsic fluorescence measurements and equilibrium studies. All spectra recorded after the 2 s deadtime can be reconstructed as linear combinations of the spectra of the initial, rapidly formed molten globule and native protein, suggesting that interconversion between the two occurs via a highly cooperative transition lacking significantly populated intermediates.

36. Oliveberg M, Tan YJ, Fersht AR: **Negative activation enthalpies in the kinetics of protein folding.** *Proc Natl Acad Sci USA* 1995, **92**:8926–8929.

Unlike many chemical reactions, protein folding tends to slow at higher (above physiological) temperatures. In this paper, data on the temperature dependence of the refolding rates of Cl2 and barnase are used to determine the relative enthalpy and heat capacity contributions to the folding transition states of these proteins. The thermodynamic properties of the major folding intermediate in barnase are also investigated.

37. Vidugiris GJA, Markley JL, Royer CA: **Evidence for a molten globule-like transition state in protein folding from determination of activation volumes.** *Biochemistry* 1995, **34**:4909–4912.

Direct detection of the activation volume for the refolding of staphylococcal nuclease from the pressure dependence of its folding rate suggests that the transition state of this process exhibits a molten-globule-like degree of compactness. The solvent-excluded expansion of this transition state contributes to the rate-limiting step in the folding of the protein.

38. Chen B-L, Baase WA, Nicholson H, Schellman JA: **Folding kinetics of T4 lysozyme and nine mutants at 12°C.** *Biochemistry* 1992, **31**:1464–1476.

39. Itzhaki LS, Evans PA, Dobson CM, Radford SE: **Tertiary interactions in the folding pathway of hen lysozyme: kinetic studies using fluorescent probes.** *Biochemistry* 1994, **33**:5212–5220.

40. Oliveberg M, Fersht AR: **Formation of electrostatic interactions on the protein-folding pathway.** *Biochemistry* 1996, **35**:2726–2737.

41. Fersht AR, Itzhaki LS, ElMasry NF, Matthews JM, Otzen DE: **Single versus parallel pathways of protein folding and fractional formation of structure in the transition state.** *Proc Natl Acad Sci USA* 1994, **91**:10426–10429.

42. Radford SE, Dobson CM: **Insights into protein folding using physical techniques: studies of lysozyme and  $\alpha$ -lactalbumin.** *Philos Trans R Soc Lond Biol* 1995, **348**:17–25.

A review of the experimental strategies used to produce the current, detailed model of the primary intermediates in the refolding of hen lysozyme. This paper provides a clear example of the power of multiple complementary probes to provide detailed insights into the nature of protein folding intermediates.

43. Plaxco KW, Spitzfaden C, Campbell ID, Dobson CM: **Rapid folding of a proline rich all  $\beta$ -sheet fibronectin type III domain.** *Proc Natl Acad Sci USA* 1996, in press.

Analysis of the folding of a fibronectin type III module by a battery of independent kinetic probes is shown to be necessary to demonstrate that the surprisingly rapid refolding of this domain reflects complete recovery of native backbone and core structure.

44. Liu X, Siegel DL, Fan P, Brodsky B, Baum J: **Direct NMR measurement of the folding kinetics of a trimeric peptide.** *Biochemistry* 1996, **35**:4306–4313.

Using peptides specifically labeled with  $^{15}\text{N}$ , the folding and association rate of trimeric collagen fibres is monitored using two-dimensional NMR with a time resolution of ~3 min. These observations suggest that the peptide folds to a significantly populated trimeric intermediate with native-like helical structure at the C terminus before folding to the native structure.

45. Fiebig KM, Schwalbe H, Buck M, Smith LJ, Dobson, CM: **Toward a description of the conformations of denatured states of proteins. Comparison of a random coil model with NMR measurements.** *J Phys Chem* 1996, **100**:2661–2666.

A model for the 'random coil' state of a polypeptide chain is developed using a Monte Carlo simulation based on an empirical sampling strategy derived from the protein structural database. While a generally good correlation between predicted and observed NOE and J-coupling constant data for lysozyme in urea suggests that the model is a reasonable representation of the distribution of structures involved, comparison of observed and predicted NOEs enables additional conformational preferences to be identified.

46. Dagget V, Li A, Itzhaki LS, Otzen DE, Fersht AR: **Structure of the transition state for folding of a protein derived from experiment and simulation.** *J Mol Biol* 1996, **257**:430–440.

Molecular dynamics simulations are performed at elevated temperatures in water to characterize the unfolding/folding transition state of Cl2. Excellent correspondence between theoretically derived and experimental values suggests that the simulated transition state provides an accurate structural model of the experimentally observed transition state.

47. Karplus M, Sali A: **Theoretical studies of protein folding and unfolding.** *Curr Opin Struct Biol* 1995, **4**:58–73.

A comprehensive review of the contribution of molecular dynamics and lattice Monte Carlo simulations to the study of protein folding.

48. Shakhnovich E, Abkevich V, Ptitsyn O: **Conserved residues and the mechanism of protein folding.** *Nature* 1996, **379**:96–98.

Lattice simulations of the folding of a series of 'homologous' sequences are used to define conserved elements in the transition state conformation of Cl2. The predicted nucleus is very similar to the folding nucleus previously defined by protein-engineering experiments.

49. Onuchic JN, Wolynes PG, Luthey-Schulten Z, Socci ND: **Toward an outline of the topography of a realistic protein folding funnel.** *Proc Natl Acad Sci USA* 1995, **92**:3626–3630.

Helix-coil transition theory and experimental data are coupled in this paper to characterize the properties of a protein folding pathway. A law of corresponding states is developed that permits comparisons to be made between experimental results and the predictions of lattice simulations of protein folding.